

Research Note

Direct measurements of black hole charge with future astrometrical missions

A.F. Zakharov^{1,2,3,4}, F. De Paolis⁵, G. Ingrosso⁵, A.A. Nucita⁵

¹ National Astronomical Observatories of Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing 100012, China

² Institute of Theoretical and Experimental Physics, 25, B.Chermushkinskaya st., Moscow, 117259, Russia,

³ Astro Space Centre of Lebedev Physics Institute, 84/32, Profsoyuznaya st., Moscow, 117810, Russia,

⁴ Joint Institute for Nuclear Research, Dubna, Russia

⁵ Department of Physics, University of Lecce and INFN, Section of Lecce, Via Arnesano, I-73100 Lecce, Italy

Received / accepted

Abstract. Recently, Zakharov et al. (2005a) considered the possibility of evaluating the spin parameter and the inclination angle for Kerr black holes in nearby galactic centers by using future advanced astrometrical instruments. A similar approach which uses the characteristic properties of gravitational retro-lensing images can be followed to measure the charge of a Reissner-Nordström black hole. Indeed, in spite of the fact that their formation might be problematic, charged black holes are objects of intensive investigations. From a theoretical point of view, it is well-known that a black hole is described by only three parameters, namely, its mass M , angular momentum J , and charge Q . Therefore, it would be important to have a method for measuring all these parameters, preferably by independent model of any. In this paper, we propose a procedure to measure the black hole charge by using the size of the retro-lensing images that can be revealed by future astrometrical missions. A discussion of the Kerr-Newmann black hole case is also offered.

Key words. black hole physics; astrometry

1. Introduction

"Black holes have no hair" means that a black hole is characterized by only three parameters ("hairs"), its mass M , angular momentum J , and charge Q (see, e.g. Misner, Thorne, and Wheeler 1973; Wald 1984 or Heusler 1998 for a more recent review). Therefore, in principle, charged black holes can be formed, although astrophysical conditions that lead to their formation may look rather problematic: see, for example, Zamir (1993); Ruffini et al. (1999, 2000); Lee, Lee & van Putten (2001); Perjes and Vasuth (2003); Ray et al. (2003); Moderski & Rogatko (2004); Vogt & Letelier (2004); Lemos & Weinberg (2004); Sereno (2004); Ghezzi & Letelier (2005). Nevertheless, one can not claim that their existence is forbidden by theoretical or observational arguments.

Charged black holes are also objects of intensive studies, since they are described by Reissner-Nordström ge-

ometry which is a static, spherically symmetrical solution of Yang-Mills-Einstein equations with fairly natural requirements on asymptotic behavior of the solutions (Gal'tsov and Ershov 1988; Gal'tsov, Volkov and Ershov 1989; Gal'tsov and Ershov 1989; Lee, Nair and Weinberg 1991). The Reissner - Nordström metric thus describes a spherically symmetric black hole with a color charge and (or) a magnetic monopole (see also Heusler 1998).

The formation of retro-lensing images (also known as mirage, shadows, or "faces" in the literature) due to the strong gravitational field effects near black holes has been investigated by several authors (Holz & Wheeler 2002; De Paolis et al. 2003, 2004; Zakharov et al. 2005a,b,c). The question that naturally arises is whether these images are observable or not. It has been shown that the retro-lensing image around the black hole at the Galactic Center (Sgr A*) due to S_2 star is observable in the K-band (peaked at $2.2 \mu\text{m}$) by the next generation infra-red space-based missions. The effects of retro-lensing image shapes due to black hole spin has also been investigated (De Paolis et al. 2004; Zakharov et al. 2005a,c,e).

Send offprint requests to: Zakharov A.F., e-mail: zakharov@itep.ru

In this paper we focus on the possibility of measuring the black hole charge as well, and we present an analytical dependence of mirage size on the black hole charge. Indeed, future space missions like Radioastron in radio band or MAXIM in X-ray band have angular resolution close to the shadow size for massive black holes in the center of our own and nearby galaxies.

2. Basic definitions and equations

The expression for the Reissner - Nordström metric in natural units ($G = c = 1$) has the form

$$ds^2 = -(1 - \frac{2M}{r} + \frac{Q^2}{r^2})dt^2 + (1 - \frac{2M}{r} + \frac{Q^2}{r^2})^{-1}dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (1)$$

Applying the Hamilton-Jacobi method to the problem of geodesics in the Reissner - Nordström metric, the motion of a test particle in the r -coordinate can be described by following equation (see, for example, Misner, Thorne, and Wheeler 1973)

$$r^4(dr/d\lambda)^2 = R(r), \quad (2)$$

where

$$\begin{aligned} R(r) &= P^2(r) - \Delta(\mu^2 r^2 + L^2), \\ P(r) &= Er^2 - eQr, \\ \Delta &= r^2 - 2Mr + Q^2. \end{aligned} \quad (3)$$

Here, the constants μ , E , L , and e are associated with the particle; i.e. μ is its mass, E is energy at infinity, L is its angular momentum at infinity, and e is the particle's charge.

We shall consider the motion of uncharged particles ($e = 0$) below. In this case, the expression for the polynomial $R(r)$ takes the form

$$R(r) = (E^2 - \mu^2)r^4 + 2M\mu r^3 - (Q^2\mu^2 + L^2)r^2 + 2ML^2r - Q^2L^2. \quad (4)$$

Depending on the multiplicities of the roots of the polynomial $R(r)$, we can have three types of motion in the r - coordinate (Zakharov 1986). In particular, by defining $r_+ = 1 + \sqrt{1 - Q^2}$, we have:

(1) if the polynomial $R(r)$ has no roots for $r \geq r_+$, a test particle is captured by the black hole;

(2) if $R(r)$ has roots and $(\partial R/\partial r)(r_{max}) \neq 0$ with $r_{max} > r_+$ (r_{max} is the maximal root), a particle is scattered after approaching the black hole;

(3) if $R(r)$ has a root and $R(r_{max}) = (\partial R/\partial r)(r_{max}) = 0$, the particle now takes an infinite proper time to approach the surface $r = const$.

If we are considering a photon ($\mu = 0$), its motion in the r -coordinate depends on the root multiplicity of the polynomial $\hat{R}(\hat{r})$

$$\hat{R}(\hat{r}) = R(r)/(M^4 E^2) = \hat{r}^4 - \xi^2 \hat{r}^2 + 2\xi^2 \hat{r} - \hat{Q}^2 \xi^2, \quad (5)$$

where $\hat{r} = r/M$, $\xi = L/(Me)$ and $\hat{Q} = Q/M$.

One can see from Eq. (5) and Eqs. (3) that the black hole charge may substantially influence the photon motion at small radii ($r \approx 1$), while the charge effect is almost negligible at large radial coordinates of photon trajectories ($r \gg 1$). In the last case, we should keep in mind that the charge may cause only small corrections to photon motion.

3. Capture cross section of photons by a Reissner - Nordström black hole

Let us consider the problem of the capture cross section of a photon by a charged black hole. It is clear that the critical value of the impact parameter for a photon to be captured by a Reissner - Nordström black hole depends on the multiplicity root condition of the polynomial $R(r)$, i. e. the condition for a vanishing discriminant (Zakharov 1988, 1991a,b, 1994a). In particular, it was shown that the vanishing discriminant condition approach is more powerful than the procedure that excludes r_{max} from the following system

$$R(r_{max}) = 0, \quad \frac{\partial R}{\partial r}(r_{max}) = 0, \quad (6)$$

as was done, for example, by Chandrasekhar (1983) to solve similar problems.

Introducing the notation $\xi^2 = l$, $Q^2 = q$, we obtain

$$R(r) = r^4 - lr^2 + 2lr - qr. \quad (7)$$

The discriminant Δ of the polynomial $R(r)$ has the form, as it was shown by Zakharov (1991a,b, 1994a):

$$\Delta = 16l^3[l^2(1 - q) + l(-8q^2 + 36q - 27) - 16q^3]. \quad (8)$$

The polynomial $R(r)$ thus has a multiple root if and only if

$$l^3[l^2(1 - q) + l(-8q^2 + 36q - 27) - 16q^3] = 0. \quad (9)$$

Excluding the case $l = 0$, which corresponds to a multiple root at $r = 0$, we find that the polynomial $R(r)$ has a multiple root for $r \geq r_+$ if and only if

$$l^2(1 - q) + l(-8q^2 + 36q - 27) - 16q^3 = 0. \quad (10)$$

If $q = 0$, we obtain the well-known result for a Schwarzschild black hole (Misner, Thorne, and Wheeler 1973; Wald 1984; Lightman et al. 1975), $l = 27$, or $L_{cr} = 3\sqrt{3}$. If $q = 1$, then $l = 16$, or $L_{cr} = 4$, which also corresponds to numerical results given by Young (1976); De Vries (2000); Takahashi (2005).

The photon capture cross section for an extremely charged black hole turns out to be considerably smaller than the capture cross section of a Schwarzschild black hole. The critical value of the impact parameter, characterizing the capture cross section for a Reissner - Nordström black hole, is determined by the equation (Zakharov 1991a,b, 1994a)

$$l = \frac{b_1 + \sqrt{b_1^2 + 64q^3(1 - q)}}{2(1 - q)}. \quad (11)$$

where $b_1 = 8q^2 - 36q + 27$.

Substituting Eq.(11) into the expression for the coefficients of the polynomial $R(r)$, it is easy to calculate the radius of the unstable circular photon orbit, which is the same as the minimum periastron distance. The orbit of a photon moving from infinity with the critical impact parameter, determined in accordance with Eq.(11), spirals into circular orbit.

As was explained by Zakharov et al. (2005a,b,c) this leads to the formation of shadows described by the critical value of L_{cr} ; or in other words, in the spherically symmetric case, shadows are circles with radii L_{cr} . Therefore, by measuring the shadow size, one could evaluate the black hole charge in black hole mass units M .




Fig. 1. Shadow (mirage) sizes are shown for selected charges of black holes $Q = 0$ (solid line), $Q = 0.5$ (short dashed line), and $Q = 1$ (long dashed line).




Fig. 2. The mirage radius l is shown as a function of the black hole charge q (l and q are given in units of M).

Table 1. The fringe sizes (in micro arc seconds) for the standard and advanced apogees B_{max} (350 000 and 3 200 000 km, respectively).

$B_{max}(\text{km}) \backslash \lambda(\text{cm})$	92	18	6.2	1.35
3.5×10^5	540	106	37	8
3.2×10^6	59	12	4	0.9

4. The space RADIOASTRON interferometer

The space-based radio telescope RADIOASTRON ¹ is planned to be launched within few next years. ² This space-based 10-meter radio telescope will be used for space – ground VLBI observations. The measurements will have extremely high angular resolutions, namely about 1 – 10 μas (in particular about 8 μas at the shortest wavelength of 1.35 cm and a standard orbit,³ and could be about 0.9 μas for the high orbit configuration at the same wavelength. Four wave bands will be used corresponding to $\lambda = 1.35$ cm, $\lambda = 6.2$ cm, $\lambda = 18$ cm, $\lambda = 92$ cm (see Table 1). A detailed calculation of the high-apogee evolving orbits (B_{max}) can be done, once the exact launch time is known.

After several years of observations, it should be possible to move the spacecraft to a much higher orbit (with apogee radius about 3.2 million km), by additional spacecraft maneuvering using the gravitational force of the Moon. The fringe sizes (in μas) for the apogee of the above-mentioned orbit and for all RADIOASTRON wavelengths are given in Table 1.

By comparing Fig. 1,2 and Table 1, one can see that there are non-negligible chances to observe such mirages around the black hole at the Galactic Center and in nearby AGNs and microquasars in the radio-band using RADIOASTRON facilities.

We also mention that this high resolution in radio band will be achieved also by Japanese VLBI project VERA (VLBI Exploration of Radio Astrometry), since the angular resolution aimed at will be at the 10 μas level (Sawad-Satoh 2000; Honma 2001). Therefore, the

¹ See web-site <http://www.asc.rssi.ru/radioastron/> for more information.

² This project was proposed by the Astro Space Center (ASC) of Lebedev Physical Institute of the Russian Academy of Sciences (RAS) in collaboration with other institutions of RAS and RosAviaKosmos. Scientists from 20 countries are developing the scientific payload for the satellite by providing by ground-based support to the mission.

³ The satellite orbit will have high apogee, and its rotation period around Earth will be 9.5 days, which evolves as a result of the weak gravitational perturbations from the Moon and the Sun. The perigee has been planned to be between 10^4 and 7×10^4 km and the apogee between 310 and 390 thousand kilometers. The basic orbit parameters will be the following: the orbital period is $P = 9.5$ days, the semi-major axis is $a = 189\,000$ km, the eccentricity is $e = 0.853$, the perigee is $H = 29\,000$ km.

only problem left is to have a powerful enough radio source to illuminate a black hole in order to have retro-lensing images detectable by such radio VLBI telescopes as RADIOASTRON or VERA.

5. Searches for mirages near Sgr A* with RADIOASTRON

Radio, near-infrared, and X-ray spectral band observations are developing very rapidly (Lo et al. 1998, 1999; Genzel et al. 2003; Ghez et al. 2004; Baganoff et al. 2001, 2003; Bower et al. 2002, 2003; Narayan 2003; Bower et al. 2004),⁴ and it is known that Sgr A* harbors the closest massive black hole with mass estimated to be $4.07 \times 10^6 M_\odot$ (Bower et al. 2004; Melia & Falcke 2001; Ghez et al. 2003; Schodel et al. 2003).

Following the idea of Falcke, Melia & Agol (2000) and of Zakharov et al. (2005a,b,c,d) we propose to use the VLBI technique to observe mirages around massive black holes and, in particular, towards the black hole at Galactic Center. To evaluate the shadow shape Falcke, Melia & Agol (2000) used the ray-tracing technique. The boundaries of the shadows are black hole mirages. We use the length parameter $r_g = \frac{GM}{c^2} = 6 \times 10^{11}$ cm for the black hole at Sgr A* and an analytical approach (see Sect. 2, 3) to calculate shadow sizes, as explained in the text. By taking the distance of Sgr A* to be $D_{GC} = 8$ kpc, the length r_g corresponds to angular size $\sim 5 \mu\text{as}$. Since the minimum arc size for the considered mirages are at least $8 r_g$, the standard RADIOASTRON resolution of about $8 \mu\text{as}$ is comparable to the required precision. The resolution in the case of the highest orbit and shortest wavelength is $\sim 1 \mu\text{as}$ (see Table 1), good enough to reconstruct the mirage shapes. As can be seen from Fig. 1 and Table 1, it is clear that, in principle, it is possible to evaluate the black hole charge Q by observing the shadow size. The mirage size difference between the extreme charged black hole and Schwarzschild black hole case is about 30% (the mirage diameter for Schwarzschild black hole is about 10.4, and for the extreme charged black hole, the diameter is equal to 8 or in black hole mass units) and typical angular sizes are about $\sim 52 \mu\text{as}$ for the Schwarzschild and $\sim 40 \mu\text{as}$ for the Reissner-Nordström black hole cases, respectively. Therefore, for Sgr A* a charged black hole could be distinguished by a Schwarzschild black hole with RADIOASTRON, at least if its charge is close to the maximal value. For stellar mass black holes, we need a much higher angular resolution to distinguish charged and uncharged black holes, since the typical shadow (mirage) angular sizes are about $2 \times 10^{-5} \mu\text{as}$, even for galactic black holes.

Actually, from the mathematical point of view we proved the following statement. For any positions of source and observer, there is an emitted photon passing close

enough to any point of the mirage that can be caught by the observer. Thus, the mirage is formed by an envelope of a family of photon trajectories or, in other words, by a caustic surface.

Mirage formation under realistic assumptions for a specific case was discussed by Falcke, Melia & Agol (2000). However, instead of solving the shadow formation problem by numerical simulations, one can use the approach developed by Zakharov (1994b, 1995); Zakharov & Repin (1999, 2002a,b, 2004a, 2003b,c,d); Zakharov et al. (2004); Zakharov (2004); Zakharov & Repin (2005a,b); Zakharov (2005). Mirages are boundaries of bright images, and shadows are also generated for accretion disks. However, for thick accretion disks as well, some parts of mirages could be observable, at least in principle.

6. Conclusions

The angular resolution of the space RADIOASTRON interferometer will be high enough to resolve radio images around black holes. By measuring the mirage shapes one should be able to evaluate the black hole mass, inclination angle (e.g. the angle between the black hole spin axis and line of sight), and spin, if the black hole distance is known. For example, for the black hole at the Galactic Center, the mirage size is $\sim 52 \mu\text{as}$ for the Schwarzschild case. In the case of a Kerr black hole (Zakharov et al. 2005a,b; De Paolis et al. 2004), the mirage is deformed depending on the black hole spin a and on the angle of the line of sight, but its size is almost the same. In the case of a Reissner-Nordström black hole, its charge changes the size of the shadows up to 30 % for the extreme charge case. Therefore, the charge of the black hole can be measured by observing the shadow size, if the other black hole parameters are known with sufficient precision. In general, one could say that a measure of the mirage shape (in size) allows to evaluate all the black hole “hairs” to be evaluated.

However, there are two kinds of difficulties when measuring mirage shapes around black holes. First, the brightness of these images or their parts (arcs) may not be sufficient for being detected by RADIOASTRON. But, numerical simulations by Falcke, Melia & Agol (2000) and Melia & Falcke (2001) give hope that the shadow brightness could be high enough to be detectable. Second, turbulent plasma effects near the black hole horizon could give essential broadening of observed images (Bower et al. 2004) leading to a confusion of the shadow image.

Recent observations of simultaneous X-ray and radio flares at 3 mm, 7 mm, 1.3 cm, and 2 cm with few-hundred second rise/fall timescales gave indirect evidence that X-ray and radio radiation from the close vicinity of Sgr A* has been detected (Baganoff et al. 2001).

A few years ago the possibility of observing the images of distant sources around black holes in the X-ray band was discussed by White (2000) and Cash et al. (2000) by using X-ray interferometer. Indeed, the aim of the

⁴ An interesting idea to use radio pulsars to investigate the region nearby black hole horizon was proposed recently by Pfahl & Loeb (2003).

MAXIM project is to realize a space-based X-ray interferometer capable of observing with angular resolution as small as $0.1\mu\text{as}$.

One could also mention that if the emitting region has a degenerate position with respect to the line of sight (for example, the inclination angle of an accretion disk is $\gtrsim 85^\circ$) strong bending effects do appear (see, Matt, Perola & Stella 1993, Zakharov & Repin 2003a).

In spite of the difficulties of measuring the shapes of shadow images, to look at black hole “faces” is an attractive challenge since mirages outline the “faces” and correspond to a fully general relativistic description of the region near the black hole horizon without any assumption about a specific model for astrophysical processes around black holes; of course, we assume that there are sources illuminating black hole surroundings. There is no doubt that the rapid growth of observational facilities will give a chance to measure the mirage shapes using not only RADIOASTRON facilities but also other instruments and spectral bands, like the X-ray interferometer MAXIM, the RADIOASTRON mission, or other space-based interferometers in millimeter and sub-millimeter bands (Miyoshi 2004).

Acknowledgements. AFZ would like to thank the Dipartimento di Fisica Università di Lecce, INFN, Sezione di Lecce and National Astronomical Observatories of Chinese Academy of Sciences, Beijing for their hospitality. AFZ thanks also the National Natural Science Foundation of China (NNSFC) for partial financial support of the work (Project # 10233050).

References

- Baganoff, F.K., et al., 2001, *Nature*, 413, 45.
 Baganoff, F.K., et al., 2003, *ApJ*, 591, 891.
 Bower, F.C., Falcke, H., Sault, R.J. & Backer, D.C., 2002, *ApJ*, 571, 843.
 Bower, F.C., Wright M.C.H., Falcke, H. & Backer, D.C., 2003, *ApJ*, 588, 331.
 Bower, F.C., Falcke, H., Herrstein, R.M., et al. 2004, *Scienceexpress*, www.scienceexpress.org/1 April 2004/.
 Cash, W., Shipley A., Osterman S., Joy M., 2000, *Nature*, 407, 160.
 Chandrasekhar, S. 1983, *Mathematical Theory of Black Holes*. Clarendon Press, Oxford.
 De Paolis, F., Geralico, A., Ingrosso, G., Nucita, A. A., 2003, *A & A*, 409, 809
 De Paolis, F., Geralico, A., Ingrosso, G., Nucita, A. A., 2004, *A & A*, 415, 1.
 Desai, K.M. & Fey A.L., 2001, *ApJS*, 133, 395.
 de Vries, A., 2000, *CQG*, 17, 123.
 Falcke, H., Melia, F., Agol, E., 2000, *ApJ*, 528, L13.
 Gal'tsov, D.V. and Ershov, A.A., 1988, *Yad. Fiz.* 4, 460.
 Gal'tsov, D.V., Volkov, M.S. and Ershov, A.A., 1989, *Tr. Inst. Fiz. Akad. Nauk Est.SSR*, 65, 160.
 Gal'tsov, D.V. and Ershov, A.A., 1989, *Phys. Lett.*, A138, 160.
 Genzel, R., R. Schödel, Ott T, et al., 2003, *Nature*, 425, 934.
 Ghez, A.M., et al., 2003, *ApJL*, 586, L127.
 Ghez, A.M., et al., 2004, *ApJL*, 601, L159.
 Ghez, A.M., et al., 2005, *ApJ*, 620, 744.
 Ghezzi, C.R., Letelier, P.S., 2005, *astro-ph/0503629*.
 Holz, D., & Wheeler J.A., 2002, *ApJ*, 578, 330.
 Honma, M. 2001, *PASJ*, 53, 233.
 Honma, M. & Kurayama T. 2002, *ApJ*, 568, 717.
 Heusler, M., 1998, *Living Reviews in Relativity*, 1998-6.
 Lee, H.K., Lee C.H., van Putten, M.H.P.M, 2001, *MNRAS*, 324, 781.
 Lee, K., Nair, V.P., Weinberg, E.J., 1991, *Fermilab Publ.*, N 91-312.
 Lemos, J.P.S., Weinberg, E. 2004, *PRD*, 69, 104004.
 Lightman, A.P., Press W.H., Price, R.H. and Teukolsky, S.A., 1975, *Problem Book in Relativity and Gravitation*, Princeton, New Jersey: Princeton University Press.
 Lo, K.Y., Shen Zh.-Q., Zhao J.-H, Ho P.T.P, 1998, *ApJL*, 508, L61.
 Lo, K.Y., Shen Zh.-Q., Zhao J.-H, Ho P.T.P, 1999. in H. Falcke, A. Cotera, W.J. Duschl, F. Melia, M.J. Rieke "The Central Parsecs of the Galaxy" (Eds.), *ASP Conference Series*, p.317.
 Matt G., Perola G.C. & Stella, L., 1993, *A&A*, 267, 2, 643.
 Melia, F., & Falcke H. 2001, *Annual Rev. Astron. & Astrophys.*, 39, 309.
 Misner, C.W., Thorne, K.S. and Wheeler, J.A. 1973 *Gravitation*, (San Francisco : Freeman)
 Miyoshi, M., 2004. *astro-ph/0412289*.
 Moderski, R., Rogatko, M.D., 2004, *PRD*, 69, 084018.
 Narayan, R., 2003, *Nature*, 425, 908.
 Perjes, Z., Vasuth M., 2003, *ApJ*, 582, 342.
 Phahl, E. & Loeb, A., 2003, *astro-ph/0309744*.
 Ray, S., Espindola, A.L., Malheiro M. et al., 2003, *PRD*, 68, 084004.
 Ruffini, R., Salmonson, J.D., Wilson J.R., Xue S.-S., 1999, *A & A*, 350, 334.
 Ruffini, R., Salmonson, J.D., Wilson J.R., Xue S.-S., 2000, *A & A*, 359, 855.
 Schödel, R., et al., 2003, *ApJ*, 596, 1015.
 Sawada-Sato, S. 2000, in H. Hirabayashi, P.G. Edwards, and D.W. Murphy (Eds.), *Proceedings of the VSOP Symposium*, held at the Institute of Space and astronomical Science, Sagamihara, Kanagawa, Japan, January 19 - 21, 2000, the Institute of Space and Astronautical Science, p.317.
 Sereno, M., 2004, *PRD*, 69, 023002.
 Takahashi, R., 2005, *astro-ph/0505316*.
 Vogt, D., & Letelier, P.S., 2004, *gr-qc/0409108*.
 Wald R.M., 1984, *General Relativity*, (The Chicago University Press)
 White, N., 2000, *Nature*, 407, 146.
 Wilkinson P.N., Narayan R., & Spencer R.E., 1994, *MNRAS*, 238, 963.
 Young, P.J., 1976, *Phys. Rev.*, D14, 3281.
 Zakharov, A.F., 1986, *Sov. Phys. JETP*, 64, 1.
 Zakharov, A.F., 1988, *Sov. Astron.*, 32, 456.
 Zakharov, A.F., 1991a, *Sov. Astron.*, 35, 147.
 Zakharov, A.F., 1991b, *Preprint ITEP*, 44-91.

- Zakharov, A.F., 1994a, CQG, 11, 1027.
- Zakharov, A.F. 1994b, MNRAS, 269, 283.
- Zakharov, A.F. 1995, In 17th Texas Symposium on Relativistic Astrophysics, Ann. NY Academy of Sciences, **759**, 550.
- Zakharov, A.F., 2004, in L. Hadzievski, T. Gvozdanov, N. Bibic (eds.), The Physics of Ionized Gases, AIP Conference Proceedings, **740**, p. 398; astro-ph/0411611.
- Zakharov, A.F., 2005, Intern. Journal Mod. Phys. A, 20, 2321.
- Zakharov, A.F., Kardashev, N.S., Lukash, V.N. & Repin, S.V. 2003, MNRAS, 342, 1325.
- Zakharov, A.F., Ma, Z. & Bao, Y., 2004, New Astronomy, 9, 663.
- Zakharov, A.F. & Repin S.V., 1999, Astronomy Reports, 43, 705.
- Zakharov, A.F. & Repin S.V., 2002a, Astronomy Reports, 46, 360.
- Zakharov, A.F., Repin S.V., 2002b, in J. Koga, T. Nakamura, K. Maeda, K. Tomita (eds.), Proceedings of the Eleven Workshop on General Relativity and Gravitation in Japan, Waseda University, Tokyo, p. 68.
- Zakharov, A.F., Repin S.V., 2003a, A & A, 406, 7.
- Zakharov, A. F., Repin, S. V., 2003b, in G. Hasinger, Th. Boller, and A.N. Parmer (eds.), XEUS - studying the evolution of the hot Universe, MPE Report 281, p. 339.
- Zakharov, A. F., Repin, S. V., 2003c, Astron. Rep., 47, 733.
- Zakharov, A.F., & Repin, S.V., 2003d, Nuovo Cimento, 118B, 1193.
- Zakharov, A.F., Repin S.V., 2004a, Advances in Space Res. 34, 1837.
- Zakharov, A.F., Repin S.V., 2004b, in L. Celniker, J. Trần Thanh Vân (eds.), Proceedings of the XIIIrd Rencontres de Blois "Frontiers of the Universe", The GIOI Publishers, p. 229.
- Zakharov A.F. & Repin S.V., 2005a, in M.S. Dimitrijević & L. Č. Popović (eds.), Proc. of the 5th Serbian Conference on Spectral Line Shapes in Astrophysics, Mem. S. A. It. (in press).
- Zakharov A.F. & Repin S.V., 2005b, New Astronomy (accepted).
- Zakharov, A.F., Nucita A.A., De Paolis F., Ingrosso G., 2005a, New Astronomy, 10, 479.
- Zakharov, A.F., Nucita A.A., De Paolis F., Ingrosso G., 2005b, in G. Vilasi, G. Esposito, G. Lambiase, G. Marmo, G. Scarpetta (eds.), Proc. of the 16th SIGRAV Conference on General Relativity and Gravitational Physics, AIP Conference Proceedings, 751, p. 227.
- Zakharov, A.F., Nucita, A.A., De Paolis, F., Ingrosso, G., 2005c, in V.A. Petrov (ed.), Proc. of the XXVII Workshop on the Fundamewntal Problems of High Energy and Field Theory, Institute for High Energy Physics, Protvino, p. 21; gr-qc/0507118.
- Zakharov, A.F., Nucita A.A., De Paolis F., Ingrosso G., 2005d, in J. Trần Thanh Vân and J. Dumarchez (eds), Proc. of XXXXth Rencontres de Moriond "Very High Energy Phenomena in the Universe" (in press).
- Zakharov, A.F., Nucita, A.A., De Paolis, F., Ingrosso, G., 2005e, in P. Chen, E. Bloom, G. Madejski, V. Petrosian (eds.), Proc. of the 22nd Texas Symposium on Relativistic Astrophysics at Stanford University, SLAC-R-752, eConf:C041213, <http://www.slac.stanford.edu/econf/C041213>, paper 1226.
- Zamir, R., 1993, ApJ, 403, 278.